

Theorem 5 *Elzaki transform amplifies the coefficients of the power series function*

$$f(t) = \sum_{n=0}^{\infty} a_n t^n, \quad (7)$$

on the new integral transform "Elzaki transform", given by

$$E[f(t)] = T(v) = \sum_{n=0}^{\infty} n! a_n v^{n+2}. \quad (8)$$

Theorem 6 *Let $f(t)$ be in A and Let $T_n(v)$ denote Elzaki transform of n th derivative, $f^{(n)}(t)$ of $f(t)$, then for $n \geq 1$,*

$$T_n(v) = \frac{T(v)}{v^n} - \sum_{k=0}^{n-1} v^{2-n+k} f^{(k)}(0). \quad (9)$$

To obtain Elzaki transform of partial derivative we use integration by parts, and then we have :

$$\begin{aligned} E\left(\frac{\partial f(x,t)}{\partial t}\right) &= \frac{1}{v}T(x,v) - v f(x,0), \\ E\left(\frac{\partial^2 f(x,t)}{\partial t^2}\right) &= \frac{1}{v^2}T(x,v) - f(x,0) - v \frac{\partial f(x,0)}{\partial t}, \end{aligned} \quad (10)$$

Properties of Elzaki transform can be found in Refs.([28],[29]), we mention only the following :

- | | |
|--------------------------|--|
| 1. $E(1) = v^2$; | 3. $E(t^n) = n!v^{n+2}$; |
| 2. $E(t) = v^3$; | 4. $E^{-1}(v^{n+2}) = \frac{t^n}{n!}$. |

2.3 Elzaki transform of fractional derivatives

To give the formula of Elzaki transform of Caputo fractional derivative, we use the Laplace transform formula for the Caputo fractional derivative [25]

$$L\{({}^c D_{0+}^{\alpha} f)(t); s\} = s^{\alpha} F(s) - \sum_{k=0}^{m-1} s^{\alpha-k-1} f^{(k)}(0),$$

where $m - 1 < \alpha \leq m$, $m \in \mathbb{N}^*$.

Theorem 7 [31] Let A defined as above. With Laplace transform $F(s)$, then the Elzaki transform $T(v)$ of $f(t)$ is given by :

$$T(v) = vF\left(\frac{1}{v}\right).$$

Theorem 8 Suppose $T(v)$ is the Elzaki transform of the function $f(t)$ then

$$E\{({}^cD_{0+}^\alpha f)(t), v\} = \frac{T(v)}{v^\alpha} - \sum_{k=0}^{n-1} v^{k-\alpha+2} f^{(k)}(0). \quad (11)$$

Proof (see [32]).

3 Modified fractional homotopy analysis transform method (MFHATM)

Kangle Wang and Sanyang Liu [15] gives the idea of the basis of this method. They consider the following general time-fractional differential equation with the initial condition as :

$$\begin{aligned} {}^cD_t^{n\alpha}U(x, t) + LU(x, t) + RU(x, t) &= g(x, t), \\ n - 1 < n\alpha \leq n, \\ U(x, 0) &= h(x), \end{aligned} \quad (12)$$

where ${}^cD_t^{n\alpha}$ is the Caputo fractional derivative operator, ${}^cD_t^{n\alpha} = \frac{\partial^{n\alpha}}{\partial t^{n\alpha}}$, L is the linear operator, R is the general nonlinear operator and $g(x, t)$ is a continuous functions.

Applying Elzaki transform on both sides of Eq.(12), we can get :

$$E[{}^cD_t^{n\alpha}U(x, t)] + E[LU(x, t) + RU(x, t) - g(x, t)] = 0, \quad (13)$$

Using the property of Elzaki transform, we have the following form:

$$E[U(x, t)] - v^{n\alpha} \sum_{k=0}^{n-1} v^{2-n\alpha+k} U^{(k)}(x, 0) + v^{n\alpha} E[LU(x, t) + RU(x, t) - g(x, t)] = 0, \quad (14)$$

Define the nonlinear operator :

$$N[\phi(x, t; p)] = E[\phi(x, t; p)] - v^{n\alpha} \sum_{k=0}^{n-1} v^{2-n\alpha+k} h^{(k)}(x, 0) \quad (15)$$

$$+ v^{n\alpha} E[L\phi(x, t; p) + R\phi(x, t; p) - g(x, t; p)]$$

By means of homotopy analysis method [1], we construct the so-called the zero-order deformation equation :

$$(1 - q)E[\phi(x, t; p) - \phi(x, t; 0)] = phH(x, t)N[\phi(x, t; p)], \quad (16)$$

where p is an embedding parameter and $p \in [0, 1]$, $H(x, t) \neq 0$ is an auxiliary function, $h \neq 0$ is an auxiliary parameter, E is an auxiliary linear Elzaki operator. When $p = 0$ and $p = 1$, we have :

$$\begin{cases} \phi(x, t; 0) = u_0(x, t), \\ \phi(x, t; 1) = u(x, t). \end{cases} \quad (17)$$

When P increases from 0 to 1, the $\phi(x, t, p)$ varies from $U_0(x, t)$ to $U(x, t)$. Expanding $\phi(x, t; p)$ in Taylor series with respect to p , we have :

$$\phi(x, t; p) = U_0(x, t) + \sum_{m=1}^{+\infty} U_m(x, t)p^m, \quad (18)$$

where

$$U_m(x, t) = \frac{1}{m!} \frac{\partial^m \phi(x, t; p)}{\partial p^m} \Big|_{p=0}. \quad (19)$$

When $p = 1$, the (18) becomes :

$$U(x, t) = U_0(x, t) + \sum_{m=1}^{+\infty} U_m(x, t). \quad (20)$$

Define the vectors :

$$\vec{U}_n = \{U_0(x, t), U_1(x, t), U_2(x, t), \dots, U_n(x, t)\}. \quad (21)$$

Differentiating (16) m -times with respect to p , then setting $p = 0$ and finally dividing them by $m!$, we obtain the so-called m th order deformation equation :

$$E[U_m(x, t) - \chi_m U_{m-1}(x, t)] = hpH(x, t)\mathfrak{R}_m(\vec{U}_{m-1}(x, t)), \quad (22)$$

where

$$\mathfrak{R}_m(\vec{U}_{m-1}(x, t)) = \frac{1}{(m-1)!} \frac{\partial^{m-1} N(x, t; p)}{\partial p^{m-1}} \Big|_{p=0}, \quad (23)$$

and

$$\chi_m = \begin{cases} 0, & m \leq 1, \\ 1, & m > 1. \end{cases}$$

Applying the inverse Elzaki transform on both sides of Eq.(22), we can obtain :

$$U_m(x, t) = \chi_m U_{m-1}(x, t) + E^{-1} \left[hpH(x, t)\mathfrak{R}_m(\vec{U}_{m-1}(x, t)) \right]. \quad (24)$$

The m th deformation equation (24) is a linear which can be easily solved. So, the solution of Eq.(12) can be written into the following form :

$$U(x, t) = \sum_{m=0}^N U_m(x, t), \quad (25)$$

when $N \rightarrow \infty$, we can obtain an accurate approximation solution of Eq.(12).

The proof of the convergence of the modified fractional homotopy analysis transform method (MFHATM) (see [2]).

4 Application of the MFHATM Method

In this section, we apply the modified fractional homotopy analysis transform method (MFHATM) for solving the following nonlinear porous medium equation with time-fractional derivative (1) in the two cases : $k = -1$ and $k = -4/3$.

4.1 Example

First, we take $k = -1$ in equation (1)[24], we get :

$${}^c D_t^\alpha u = (u^{-1}u_x)_x, \quad 0 < \alpha \leq 1, \quad (26)$$

with the initial condition :

$$u(x, 0) = \frac{1}{x}. \quad (27)$$

Applying Elzaki transform on both sides of Eq.(26), we can get :

$$E[u] - v^2 u(x, 0) = v^\alpha E[(u^{-1}u_x)_x] \quad (28)$$

From (28) and the initial condition (27), we have :

$$E[u] - v^2 \frac{1}{x} - v^\alpha E[(u^{-1}u_x)_x] = 0. \quad (29)$$

We take the nonlinear part as :

$$N[\phi(x, t, p)] = E[\phi] - v^2 \frac{1}{x} - v^\alpha E[(\phi^{-1}\phi_x)_x]. \quad (30)$$

We construct the so-called the zero-order deformation equation with assumption $H(x; t) = 1$, we have :

$$(1 - q)E[\phi(x, t; p) - \phi(x, t; 0)] = phN[\phi(x, t; p)]. \quad (31)$$

When $p = 0$ and $p = 1$, we can obtain :

$$\begin{cases} \phi(x, t; 0) = u_0(x, t), \\ \phi(x, t; 1) = u(x, t). \end{cases}$$

Therefore, we have the m th order deformation equation :

$$E[u_m(x, t) - \chi_m u_{m-1}(x, t)] = h\mathfrak{R}_m(\vec{u}_{m-1}(x, t)). \quad (32)$$

Operating the inverse Elzaki operator on both sides of Eq.(32), we get :

$$u_m(x, t) = \chi_m u_{m-1}(x, t) + E^{-1}[h\mathfrak{R}_m(\vec{u}_{m-1}(x, t))]. \quad (33)$$

From Eq.(33), we have :

$$\begin{aligned} u_1(x, t) &= hE^{-1}[\mathfrak{R}_1(\vec{u}_0(x, t))], \\ u_2(x, t) &= u_1 + hE^{-1}[\mathfrak{R}_2(\vec{u}_1(x, t))], \\ u_3(x, t) &= u_2 + hE^{-1}[\mathfrak{R}_3(\vec{u}_2(x, t))], \\ &\vdots \end{aligned} \tag{34}$$

where

$$\begin{aligned} \mathfrak{R}_1(\vec{u}_0(x, t)) &= E[u_0] - v^2 \frac{1}{x} - v^\alpha E[(u_0^{-1}u_{0x})_x], \\ \mathfrak{R}_2(\vec{u}_1(x, t)) &= E[u_1] - v^\alpha E[(u_0^{-1}u_{1x} - u_0^{-2}u_1u_{0x})_x], \\ \mathfrak{R}_3(\vec{u}_2(x, t)) &= E[u_2] - v^\alpha E[(u_0^{-3}u_1^2u_{0x} - u_0^{-2}u_2u_{0x} - u_0^{-2}u_1u_{1x} + u_0^{-1}u_{2x})_x], \\ &\vdots \end{aligned} \tag{35}$$

Using the initial condition (27), the iteration formulas (34) and (35), we obtain :

$$\begin{aligned} u_0(x, t) &= \frac{1}{x}, \\ u_1(x, t) &= -\frac{h}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)}, \\ u_2(x, t) &= -\frac{h}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{h^2}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} + 2\frac{h^2}{x^3} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}, \\ u_3(x, t) &= -\frac{h}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{h^2}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} + 4\frac{h^2}{x^3} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\ &\quad - \frac{h^2}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{h^3}{x^2} \frac{t^\alpha}{\Gamma(\alpha + 1)} + 4\frac{h^3}{x^3} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\ &\quad + 3\frac{h^3}{x^4} \frac{\Gamma(2\alpha + 1)}{\Gamma(\alpha + 1)^2} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} - 12\frac{h^3}{x^4} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} \\ &\quad \vdots \end{aligned} \tag{36}$$

Thus, we use four terms in evaluating the approximate solution :

$$u(x, t) = \sum_{m=0}^3 u_m(x, t).$$

When $h = -1$, the approximate solution of Eq.(26), is given by :

$$\begin{aligned} u(x, t) &= \sum_{m=0}^3 u_m(x, t) \\ &= \frac{1}{x} + \frac{1}{\Gamma(\alpha + 1)} \frac{t^\alpha}{x^2} + \frac{2}{\Gamma(2\alpha + 1)} \frac{t^{2\alpha}}{x^3} - \frac{3\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2\Gamma(3\alpha + 1)} \frac{t^{3\alpha}}{x^4} \\ &\quad + \frac{12}{\Gamma(3\alpha + 1)} \frac{t^{3\alpha}}{x^4}. \end{aligned}$$


If $\alpha = 1$, we obtain :

$$u(x, t) = \frac{1}{x} + \frac{t}{x^2} + \frac{t^2}{x^3} + \frac{t^3}{x^4} + \dots$$


That gives :

$$u(x, t) = \frac{1}{x - t}, \quad \left| \frac{t}{x} \right| < 1, \quad x \neq 0,$$


which is an exact solution to the porous medium equation as presented in [24].



(a)



(b)



(c)

Fig. 1 : (a) Exact solution, (b) the approximate solution in the case $\alpha = 1$, (c) The exact solution and approximate solutions to the Eq.(26) for different values of α when $x = 2$.

4.2 Example

Second, we take $k = -4/3$ in equation (1)[24], we get :

$${}^c D_t^\alpha u = (u^{-4/3} u_x)_x, \quad 0 < \alpha \leq 1, \quad (37)$$

with the initial condition :

$$u(x, 0) = (2x)^{-3/4}. \quad (38)$$

Applying Elzaki transform on both sides of Eq.(37), we get :

$$E[u] - v^2 u(x, 0) = v^\alpha E[(u^{-4/3} u_x)_x] \quad (39)$$

From (39) and the initial condition (38), we have :

$$E[u] - v^2 (2x)^{-3/4} - v^\alpha E[(u^{-4/3} u_x)_x] = 0.$$

We take the nonlinear part as :

$$N[\phi(x, t, p)] = E[\phi] - v^2 (2x)^{-3/4} - v^\alpha E\left[\left(\phi^{-4/3} \phi_x\right)_x\right].$$

We construct the so-called the zero-order deformation equation with assumption $H(x; t) = 1$, we have :

$$(1 - q)E[\phi(x, t; p) - \phi(x, t; 0) = phN[\phi(x, t; p)].$$

When $p = 0$ and $p = 1$, we can obtain :

$$\begin{cases} \phi(x, t; 0) = u_0(x, t), \\ \phi(x, t; 1) = u(x, t). \end{cases}$$

Therefore, we have the m th order deformation equation :

$$E[u_m(x, t) - \chi_m u_{m-1}(x, t)] = h\mathfrak{R}_m(\vec{u}_{m-1}(x, t)). \quad (40)$$

Operating the inverse Elzaki operator on both sides of Eq.(40), we get :

$$u_m(x, t) = \chi_m u_{m-1}(x, t) + E^{-1}[h\mathfrak{R}_m(\vec{u}_{m-1}(x, t))], \quad (41)$$

From Eq.(41), we have :

$$\begin{aligned} u_1(x, t) &= hE^{-1}[\mathfrak{R}_1(\vec{u}_0(x, t))], \\ u_2(x, t) &= u_1 + hE^{-1}[\mathfrak{R}_2(\vec{u}_1(x, t))], \\ u_3(x, t) &= u_2 + hE^{-1}[\mathfrak{R}_3(\vec{u}_2(x, t))], \\ &\vdots \end{aligned} \quad (42)$$

where

$$\begin{aligned} \mathfrak{R}_1(\vec{u}_0(x, t)) &= E[u_0] - v^2(2x)^{-3/4} - v^\alpha E \left[\left(u_0^{-4/3} u_{0x} \right)_x \right], \\ \mathfrak{R}_2(\vec{u}_1(x, t)) &= E[u_1] - v^\alpha E \left[\left(-\frac{4}{3} u_0^{-7/3} u_1 u_{0x} + u_0^{-4/3} u_{1x} \right)_x \right], \\ \mathfrak{R}_3(\vec{u}_2(x, t)) &= E[u_2] - v^\alpha E \left[\left(\frac{14}{9} u_0^{-10/3} u_1^2 u_{0x} - \frac{4}{3} u_0^{-7/3} u_2 u_{0x} \right. \right. \\ &\quad \left. \left. - \frac{4}{3} u_0^{-7/3} u_1 u_{1x} + u_0^{-4/3} u_{2x} \right)_x \right], \\ &\vdots \end{aligned} \quad (43)$$

Using the initial condition (38), the iteration formulas (42) and (43), we obtain :

$$\begin{aligned}
 u_0(x, t) &= (2x)^{-3/4}, \\
 u_1(x, t) &= -\frac{9}{4}h(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)}, \\
 u_2(x, t) &= -\frac{9}{4}h(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{9}{4}h^2(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} + \frac{189}{16}h^2(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}, \\
 u_3(x, t) &= -\frac{9}{4}h(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{9}{4}h^2(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} + \frac{189}{16}h^2(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 &\quad - \frac{9}{4}h^2(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{9}{4}h^3(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} + \frac{189}{16}h^3(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 &\quad + \frac{1089}{32}h^3(2x)^{-15/4} \frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} + \frac{189}{16}h^2(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 &\quad + \frac{189}{16}h^3(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} - \frac{14553}{64}h^3(2x)^{-15/4} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} \\
 &\quad + \frac{2079}{16}h^2(2x)^{-15/4} \frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)}. \\
 &\quad \vdots
 \end{aligned}$$

Thus, we use four terms in evaluating the approximate solution :

$$u(x, t) = \sum_{m=0}^3 u_m(x, t).$$

When $h = -1$, the approximate solution of Eq.(37), is given by :

$$\begin{aligned}
 u(x, t) &= \sum_{m=0}^3 u_m(x, t) \\
 &= (2x)^{-3/4} + \frac{9}{4}(2x)^{-7/4} \frac{t^\alpha}{\Gamma(\alpha + 1)} + \frac{189}{16}(2x)^{-11/4} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 &\quad + \frac{3069}{32}(2x)^{-15/4} \frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)} + \frac{14553}{64}(2x)^{-15/4} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)}.
 \end{aligned}$$

If $\alpha = 1$, we obtain :

$$u(x, t) = (2x)^{-3/4} + \frac{9}{4}(2x)^{-7/4} t + \frac{189}{32}(2x)^{-11/4} t^2 + \frac{8943}{128}(2x)^{-15/4} + \dots ,$$

or

$$u(x, t) = 2^{-3/4} \times x^{-3/4} + (9 \times 2^{-15/4} \times x^{-7/4})t + (189 \times 2^{-31/4} \times x^{-11/4})t^2 + (8943 \times 2^{-43/4} \times x^{-15/4})t^3 + \dots,$$

which is the same approximate solution to the porous medium equation as presented in [24].




Fig. 2 : (a) The approximate solution when $\alpha = 1$, (b) The approximate solution when $\alpha = 0.98$, (c) The approximate solutions to the Eq.(37) for different values of α when $x = 0.2$.

Remark 9 For graph approximate solutions, we took only four terms.

5 Conclusion

The coupling of homotopy analysis method (HAM) and Elzaki transform method proved very effective to solve nonlinear partial differential equations. The modified fractional homotopy analysis transform method (MFHATM), is suitable for such problems and is very user friendly. From the obtained results, it is clear that the MFHATM yields very accurate, exact and approximate solutions using only a few iterates. As a result, the conclusion that comes through this work is that MFHATM can be applied to other nonlinear fractional partial differential equations of higher order, due to the efficiency and flexibility in the application to get the possible results.

References

- [1] S.J. Liao, "*The proposed homotopy analysis technique for the solution of nonlinear problems*", Ph.D. Thesis, Shanghai Jiao Tong University, (1992).
- [2] S.J. Liao, "*Beyond Perturbation: Introduction to Homotopy Analysis Method*", Chapman and Hall/CRC Press, Boca Raton, (2003).
- [3] S.J. Liao, "*On the homotopy analysis method for nonlinear problems*", Appl. Math. Comput., 147, (2004), 499-513.
- [4] G. Adomian, "*Nonlinear Stochastic Systems Theory and Applications to Physics*", Kluwer Academic Publishers, Netherlands, (1989).
- [5] G. Adomian, R. Rach, "*Equality of partial solutions in the decomposition method for linear or nonlinear partial differential equations*", Comput. Math. Appl., 10, (1990), 9-12.
- [6] G. Adomian, "*Solution of physical problems by decomposition*", Comput. Math. Appl. 27, (1994), 145-154.
- [7] J.H. He, "*A new approach to nonlinear partial differential equations*", Comm. Nonlinear Sci. Numer. Simul., 2, (1997), 203-205.

- [8] J.H. He, "Approximate analytical solution for seepage flow with fractional derivatives in porous media", *Comput. Meth. Appl. Mech. Eng.*, 167, (1998), 57-68.
- [9] J.H. He, "A variational iteration approach to nonlinear problems and its applications", *Mech. Appl.*, 20, (1998), 30-31.
- [10] J.H. He, "Homotopy perturbation technique", *Comput. Meth. Appl. Mech. Eng.*, 178, (1999), 257-262.
- [11] J.H. He, "A coupling method of homotopy technique and perturbation technique for nonlinear problems", *Int. J. Nonlinear Mech.*, 35, (2000), 37-43.
- [12] J.H. He, "A new perturbation technique which is also valid for large parameters", *J. Sound Vib.*, 229, (2000), 1257-1263.
- [13] M. Zurigat, "Solving Fractional Oscillators Using Laplace Homotopy Analysis Method", *Annals of the University of Craiova, Math. Comp. Sci. Series*, 38, (2011), 1-11.
- [14] S. Rathore, D. Kumar, J. Singh, S. Gupta, "Homotopy Analysis Sumudu Transform Method for Nonlinear Equations", *Int. J. Industrial Math.*, 4, (2012), 301-314.
- [15] K. Wang, S. Liu, "Application of new iterative transform method and modified fractional homotopy analysis transform method for fractional Fornberg-Whitham equation", *J. Nonlinear Sci. Appl.* 9 (2016), 2419-2433.
- [16] S. A. Khuri, "A Laplace decomposition algorithm applied to a class of nonlinear differential equations", *J. Math. Annl. Appl.*, 4, (2001), 141-155.
- [17] D. Kumar, J. Singh, S. Rathore, "Sumudu Decomposition Method for Nonlinear Equations", *Int. Math. For.*, 7, (2012), 515 - 521.
- [18] M. Khalid, M. Sultana, F. Zaidi, U. Arshad, "An Elzaki Transform Decomposition Algorithm Applied to a Class of Non-Linear Differential Equations", *J. of Natural Sci. Res.*, 5, (2015), 48-55.

- [19] A.S. Arife, A. Yildirim, "New Modified Variational Iteration Transform Method (MVITM) for solving eighth-order Boundary value problems in one Step", W. Appl. Sci. J., 13, (2011), 2186 -2190.
- [20] A.S. Abedl-Rady, S.Z. Rida, A.A.M. Arafa, H.R. Abedl-Rahim, "Variational Iteration Sumudu Transform Method for Solving Fractional Nonlinear Gas Dynamics Equation", Int. J. Res. Stu. Sci. Eng. Tech., 1, (2014), 82-90.
- [21] J. Manafian, I. Zamanpour, "Application of the ADM Elzaki and VIM Elzaki transform for solving the nonlinear partial differential equations", Sci. Road J., 2, (2014), 37-50.
- [22] S. Kumara, A. Yildirim, Y. Khan, L. Weid, "A fractional model of the diffusion equation and its analytical solution using Laplace transform", Scientia Iranica B 19, (2012), 1117-1123.
- [23] J. Singh, D. Kumar, Sushila, "Homotopy Perturbation Sumudu Transform Method for Nonlinear Equations", Adv. Theor. Appl. Mech., 4, (2011), 165 - 175.
- [24] P.K.G. Bhadane, V.H. Pradhan, "Elzaki Trnsform Homotopy Perturbation Method for Solving Porous Medium Equation", Int. J. Res. Eng. Tech. 2(12), (2013), 116-119.
- [25] I. Podlubny, "Fractional Differential Equations", Academic Press, San Diego, CA, (1999).
- [26] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, "Theory and Applications of Fractional Differential Equations", Elsevier, Amsterdam (2006).
- [27] K. Diethelm, "The Analysis Fractional Differential Equations", Springer-Verlag Berlin Heidelberg (2010).
- [28] T.M. Elzaki, S.M. Elzaki, E.A. Elnour, "On the New Integral Transform "ELzaki Transform" Fundamental Properties Investigations and Applications", Glo. J. Math. Sci., 4, (2012), 1-13.
- [29] T.M. Elzaki, E.M.A. Hilal, "Homotopy Perturbation and Elzaki Transform for Solving Nonlinear Partial Differential Equations", Math. Theor. Mod., 2, (2012), 33-42.

- [30] E.M. Abd Elmohmoud, T.M. Elzaki, "*Elzaki Transform of Derivative Expressed by Heaviside Function*", W. Appl. Sci. J., 32, (2014), 1686-1689.
- [31] T.M. Elzaki, S.M. Ezaki; "*On the Connections Between Laplace and ELzaki Transforms*", Adv. Theo. Appl. Math. 6, (1), (2011), 1–10.
- [32] A. Neamaty, B. Agheli, R. Darzi; "*New Integral Transform for Solving Nonlinear Partial Differential Equations of fractional order*", Theo. Appr. Appl. 10, (1), (2014), 69-86.